

Graviton KK resonant mode in the correction of the Newton's law from 6D braneworlds

J. C. B. Araújo,^{*} D. F. S. Veras,[†] D. M. Dantas,[‡] and C. A. S. Almeida[§]

*Departamento de Física, Universidade Federal do Ceará (UFC),
Campus do Pici, Caixa Postal 6030, 60455-760, Fortaleza, Ceará, Brazil*

Abstract

In this work, we derive an expression for the correction in the Newton's law of gravitation due to the gravitational Kaluza-Klein states in a general thick string-like braneworld scenario in six dimensions. In order to analyze corrections to Newton's law we study the gravity fluctuations in a 3-brane placed in a transverse resolved conifold and use suitable numerical methods to attain the massive spectrum and the corresponding eigenfunctions. Such braneworld model has a resolution parameter which removes the conical singularity. The correction has an exponentially suppressed mass term and depends on the values of the eigenfunctions and warp factors computed at the core peak of the brane. The spectrum is real and monotonically increased, as desired. However, the resolution parameter must assume moderate values to have physically acceptable states. Moreover, the trapped massless mode regains the 4D gravity and it is displaced from the origin, sharing similar profile with the energy density of brane for small values of resolution parameter. Finally, for the singular conifold, we found that a non-first eigenstate is a resonant mode. Such excited state is the largest contributor to corrections in the Newtonian potential.

Keywords: Braneworlds, Gravity Localization, Resolved Conifold, Newton's law correction

^{*} julio@fisica.ufc.br

[†] franklin@fisica.ufc.br

[‡] davi@fisica.ufc.br

[§] carlos@fisica.ufc.br

I. INTRODUCTION

The braneworld hypothesis has changed the way we understand the Universe introducing the idea that it may be thought as a hypersurface embedded in a higher dimensional bulk space-time. The braneworld concept emerged from string theory [1] and unification models [2] and yields good explanations to several fundamental phenomena in High Energy Physics such as the hierarchy problem [3, 4], the dark matter origin [5], the small value of the cosmological constant [6] and the cosmic acceleration [7].

The seminal papers of Lisa Randall and Raman Sundrum (RS) [3, 8] assume an Anti-de Sitter spacetime (AdS_5) through a warped product between a 3-brane and a single extra dimension. As a result, the hierarchy problem can be explained by means of an exponential decreasing factor between the Planck scale and the weak scale along the extra dimension [3]. Furthermore, the bulk massive gravitons are responsible for a small correction to the Newtonian potential of order $\mathcal{O}(r^{-2})$ [8]. Thenceforth, several works were carried out enhancing the RS models, such as proving the stability of the geometrical solution [9], providing a physical source for the geometry [10–16], allowing the Standard Model fields to propagate in the bulk [17–19], and extending the model to higher dimensions [20–23].

The extension of the RS models to six dimensions with an axial symmetry was proposed by Gherghetta and Shaposhnikov (GS) in the Ref. [23], wherein the transverse manifold is a circle. In the AdS_6 GS *string-like scenario*, the problem of mass hierarchy is solved without any requirement of fine tuning between the bulk cosmological constant and the brane tension [23]. Furthermore, the Kaluza-Klein (KK) excitations generate corrections to the Newtonian potential of order $\mathcal{O}(r^{-3})$. Moreover, the string-like models has the advantage of trapping free gauge fields [24] and minimally coupled Dirac fermions [25]. Besides, in the Lorentz invariance violation context, the string-like defect with a bulk dependent cosmological constant can yield a massless four-dimensional graviton [26], which is not possible in the thin 5D model [27].

Although the thin string-like model has prominent advantages over the domain-wall models, it does not satisfy all the regularity conditions inside the core of the defect and the dominant energy condition as well [28]. However, these issues are overcome in the thick string-like models [29–33]. The transverse manifold possesses its own internal symmetries, which exert influences on the geometrical and physical properties of the braneworld model [32, 34, 35]. Moreover, the $6D$ warped models have other interesting features. For instance, Refs. [35, 36] suggest defects with two angular extra dimensions, wherein the angular momentum in the transverse space is correlated

to the three generations of fundamental $4D$ fermions. Additionally, Ref. [37] uses a single fermion family in $6D$ to explain the mass hierarchy of neutrinos.

An interesting six dimensional braneworld model with axial symmetry, proposed by Silva and Almeida, uses a section of the resolved conifold (RC) as transverse space and traps the scalar [32] and matter fields [38, 39]. The resulting scenario has a parameter a which controls the singularity on the tip of the cone (see Ref. [38] and references therein). Moreover, the geometric flow provided by the resolution parameter changes the properties of the analogue quantum potential for the KK massives states. An important issue in the thick string-like braneworld scenarios concerns with the core of the brane, which has its maximum displaced from the origin [29, 31–33]. Such quality is due to the influence of the geometric flow in the physics of the brane [31].

In this letter, we are interested in study the gravity localization on a 3-brane placed in a transverse warped resolved conifold. We derive a general expression for the correction in the Newtonian gravitation potential due to the graviton KK states in a thick string-like braneworld scenario. Furthermore, we use suitable numerical techniques to attain the graviton spectrum and the corresponding eigenfunctions. With the eigensolutions in hands, we were able to analyse the influence of the parameters of the model in the correction of the gravitational potential.

II. BULK GEOMETRY AND PHYSICAL PROPERTIES

In this section, we will present a brief review of the most important geometric and physical properties of the resolved conifold (RC) braneworld model. First of all, the action for a $6D$ spacetime can be denoted as [24]

$$S_6 = \int_{\mathcal{M}_6} \left(\frac{1}{2\kappa_6} R - \Lambda + \mathcal{L}_m \right) \sqrt{-g} d^6x . \quad (1)$$

Here, $\kappa_6 = 8\pi G_N = 8\pi/M_6^4$, where G_N is the gravitational constant and M_6 is the six-dimensional bulk Planck mass. Moreover, $\mathcal{L}_m$ is the matter Lagrangian for the source of the geometry and the bulk cosmological constant Λ has dimension $[\Lambda] = L^{-6} = M^6$. From this action, the $6D$ Einstein equations are obtained as follows:

$$R_{AB} - \frac{R}{2} g_{AB} = -\kappa_6 (\Lambda g_{AB} + T_{AB}) . \quad (2)$$

Now, consider a static and axisymmetric warped metric between the 3-brane $\mathcal{M}_4$ and the transverse space given by [23]

$$ds_6^2 = \sigma(\rho) \eta_{\mu\nu}(x^\zeta) dx^\mu dx^\nu + d\rho^2 + \gamma(\rho) d\theta^2 , \quad (3)$$

where $\sigma(\rho)$ is the so-called warp function, x^ζ are on-brane coordinates and (ρ, θ) are coordinates of the transverse manifold. The metric has axial symmetry, so that, $r \in [0, \infty)$ and $\theta \in [0, 2\pi]$. The function $\gamma(\rho)$ is an angular warp factor with dimension L^2 . Besides, we adopt the sign convention for the Minkowski metric as $\eta_{\mu\nu} = \text{diag}(-, +, +, +)$.

Furthermore, an axisymmetric ansatz for the energy-momentum tensor may be considered as $T_N^M = T_M \delta_N^M$ [23, 24], which together the metric ansatz (3), leads the energy density to the form [23, 24, 31]

$$t_0(r) = -\frac{1}{\kappa_6} \left[\frac{3}{2} \left(\frac{\sigma'}{\sigma} \right)' + \frac{3}{2} \left(\frac{\sigma'}{\sigma} \right)^2 + \frac{1}{2} \left(\frac{\gamma'}{\gamma} \right)' - \frac{1}{4} \left(\frac{\gamma'}{\gamma} \right)^2 + \frac{3}{4} \frac{\sigma'}{\sigma} \frac{\gamma'}{\gamma} \right] - \Lambda . \quad (4)$$

In order to obtain a regular geometry free of singularities, we need to impose the following regularity conditions at the origin [28, 40]:

$$\sigma(\rho)|_{\rho=0} = \text{cte} , \quad \sigma'(\rho)|_{\rho=0} = 0 , \quad (5)$$

$$\gamma(\rho)|_{\rho=0} = 0 , \quad \left(\sqrt{\gamma(\rho)} \right)' \Big|_{\rho=0} = 1 . \quad (6)$$

In the Gherghetta-Shaposhnikov (GS) thin string-like model, the solutions of the warp factor outside the core of the brane is obtained as [23]:

$$\sigma(\rho) = e^{-c\rho} , \quad \gamma(\rho) = R_0^2 \sigma(\rho) . \quad (7)$$

The parameter $c^2 = -\frac{2}{5}\kappa_6\Lambda$ can be obtained from the vacuum solution of the Eq. (4). Note that only the first condition in the Eq. (6) for the GS model is complied. Moreover, the scalar curvature yields to a pure AdS_6 spacetime where $R = -\frac{15}{2}c^2$ [23].

In the RC braneworld, instead use of the disc, it is chosen a 2-section of the resolved conifold as the transverse manifold [32, 41, 42]. The following warp factors are considered:

$$\sigma(\rho) = e^{-(c\rho - \tanh c\rho)} , \quad \gamma(\rho) = \left(\frac{u(\rho, a)^2 + 6a^2}{6} \right) \sigma(\rho) , \quad (8)$$

where a is the resolution parameter and the function u has the form [38, 39]

$$u_a(\rho) = \begin{cases} \rho , & a = 0 \\ -i\sqrt{6}aE\left(\arcsin\left(\frac{i\rho}{3a}\right), \frac{2}{3}\right) , & a \neq 0 \end{cases} \quad (9)$$

where E represents the elliptic integral of the second kind.

We plot in the Fig. 1 the warp factors $\sigma(\rho)$ and $\gamma(\rho)$ for $c = 1.0$ and different values of a . The thin string case is embedded. The warp function $\sigma(\rho)$ was first proposed in the *string-cigar*




Figure 1. Warp factors $\sigma(\rho)$ and $\gamma(\rho)$ for $c = 1.0$.




Figure 2. Energy Density $t_0(\rho)$ for different values of a .

braneworld [31]. Note that it recovers the usual thin string-like exponential behaviour asymptotically. Moreover, differently from the thin string-like models, the warp function satisfies all the regularity conditions at the origin. Therefore, models with this warp function can be realized as a near brane correction to the thin string-like models [31, 33].

The angular ansatz given in the Eq. (8) presents noteworthy features at the origin. Note that $\gamma(0) = a^2$, thus the resolution parameter removes the conical singularity at the origin [31]. Furthermore, for $\rho = 0$, the 4-brane is a $\mathcal{M}_5$ whose metric is $ds_5^2 = \eta_{\mu\nu}(x^\zeta)dx^\mu dx^\nu + a^2 d\theta^2$. Therefore, the geometric flow of the resolved conifold, driven by the resolution parameter, leads to a dimensional reduction $\mathcal{M}_6 \rightarrow \mathcal{M}_5$. The string-like dimensional reduction $\mathcal{M}_6 \rightarrow \mathcal{M}_4$ is achieved for $a = 0$. Such features are of substantial relevance in the localization of the spin-1 and spin-1/2 fields [38, 39].

Due to the intricate form of the Eq. (4) for the warp factors given by Eqs. (8), we present the plot of the energy density in the Fig. 2 for different values of a . Note that the maximum of the energy density is shifted from the origin. Such feature is frequently observed in thick string-like braneworld models [29, 31, 33]. This displacement in the core peak of the thick branes has important influences in the massive modes of the gravitational [43], gauge [44], fermionic [45, 46] and exotic elko [47] fields. Note from the Fig. 2 that the position of the maximum of the energy density has not great variations with the resolution parameter. Despite of this particular feature, the resolution parameter is of great importance in the mass spectrum, which has a direct influence on corrections to the gravitational potential as we will present in details in the Sec. V.

III. METRIC PERTURBATIONS

We will now study the gravity fluctuations on a 3-brane placed in a transverse resolved conifold (RC). It was shown recently that the scalar, gauge and fermionic fields are trapped in such braneworld model [32, 38, 39]. Besides, the study of gravity localization is quite relevant since it provides phenomenology implications [8, 10]. This analysis will be accomplished in the Sec. IV through corrections in the Newton's law.

First of all, consider a small perturbation $h_{\mu\nu}$ in the background metric in the form

$$ds_6^2 = \sigma(\rho; c)(\eta_{\mu\nu} + h_{\mu\nu})dx^\mu dx^\nu + d\rho^2 + \gamma(\rho; c, a)d\theta^2 . \quad (10)$$

Imposing the transverse-traceless gauge $\nabla^\mu h_{\mu\nu} = 0$, the linearization of the Einstein equations yields to the following equation for the gravitational perturbation [23, 29, 31]:

$$\partial_A(\sqrt{-g_6} g^{AB} \partial_B h_{\mu\nu}) = 0 . \quad (11)$$

Performing the well-known Kaluza-Klein decomposition

$$h_{\mu\nu}(x^\zeta, \rho, \theta) = \tilde{h}_{\mu\nu}(x^\zeta) \sum_{n=0}^{\infty} \phi_n(\rho) e^{il\theta} , \quad (12)$$

where n and l are integers, and imposing the mass condition $\square_4 \tilde{h}_{\mu\nu}(x^\zeta) = m_n^2 \tilde{h}_{\mu\nu}(x^\zeta)$, the radial modes satisfy the equation

$$\phi''(\rho) + \left(2\frac{\sigma'}{\sigma} + \frac{1}{2}\frac{\gamma'}{\gamma} \right) \phi'(\rho) + M^2(\rho)\sigma(\rho)\phi(\rho) = 0 , \quad (13)$$

where the primes denote derivatives with respect to ρ . The factor $M^2(\rho) = \left(m^2 - \frac{\sigma(\rho)}{\gamma(\rho)} l^2 \right)$ is an effective mass containing orbital angular momentum contributions l .

Due to the axial symmetry and in order to guarantee the self-adjointness of the differential operator in the Eq. (11), we adopt the following boundary conditions on ϕ_m [23, 24, 31, 33]:

$$\phi'_n(0) = \phi'_n(\infty) = 0 . \quad (14)$$

Furthermore, these modes satisfy the following orthogonality condition:

$$\int_0^\infty \sigma(\rho) \sqrt{\gamma(\rho)} [\phi_{n_1}^*(\rho) \phi_{n_2}(\rho)] d\rho = \delta_{n_1 n_2} . \quad (15)$$

For $M^2(\rho) = 0$ (gravitational massless mode), a solution for the Eq. (13) satisfying the boundary conditions (14) is a constant $\phi = \phi_0$ [23]. Hence, from the orthonormality condition (15), we have:

$$\phi_0 = \left(\int_0^\infty \sigma(\rho) \sqrt{\gamma(\rho)} d\rho \right)^{-\frac{1}{2}} . \quad (16)$$

Since that for both GS and RC models the warp factors are exponentially decreasing, the constant solutions are finite and hence the zero mode is localized. For the GS model, the solution is $\phi_0 = \sqrt{\frac{3c}{2R_0}}$.

For $M^2(\rho) \neq 0$ (massive modes), the differential equation (13) with the warp factors given by Eq. (8), becomes

$$\phi''(\rho) - \left(\frac{5}{2}c \tanh^2(c\rho) - \frac{1}{2} \frac{u(u^2 + 9a^2)^{\frac{1}{2}}}{(u^2 + 6a^2)^{\frac{3}{2}}} \right) \phi'(\rho) + \left(m^2 - \frac{\sigma(\rho)}{\gamma(\rho)} l^2 \right) e^{(c\rho - \tanh c\rho)} \phi(\rho) = 0 . \quad (17)$$

In the limit $\rho \rightarrow \infty$, we have

$$\phi''(\rho) - \frac{5}{2}c \phi'(\rho) + m^2 e^{(c\rho)} \phi(\rho) = 0 , \quad (18)$$

which present the same form of the GS model [23] for a re-scaled mass $m^2 \rightarrow m^2/e$. Thus, the thin string-like behaviour is recovered asymptotically, as expected from thick string-like braneworlds [31, 33].

Note further that the angular eigenvalue l induces a degeneracy in the mass spectrum. Thus, hereinafter we will deal with s -wave solutions. The s -wave solution in the thin limit has the analytical solution [23]

$$\phi_m(\rho) = e^{(5/4)c\rho} \left[C_1 J_{5/2} \left(\frac{2m}{c} e^{\frac{c\rho}{2}} \right) + C_2 Y_{5/2} \left(\frac{2m}{c} e^{\frac{c\rho}{2}} \right) \right] , \quad (19)$$

which diverges. Hence, there are not massive bound states in the resolved conifold braneworld model, as well.

IV. CORRECTIONS IN THE NEWTONIAN POTENTIAL

Another approach in the study of the gravitational massive modes consists in turn the Kaluza-Klein equation (13) into a Schrödinger-like one. Such procedure provides the study of the phenomenological implications of the braneworld hypothesis via corrections in the Newton's law of gravitation [8, 10]. Here, for the first time, it is obtained an expression for a general thick braneworld scenario in order to calculate corrections in the Newton's law.

First of all, the transformation of coordinate $z(\rho) = \int_0^\rho \sigma^{-\frac{1}{2}}(\rho') d\rho'$ provides a conformally plane metric as $ds_6^2 = \sigma(z)(\eta_{\mu\nu} dx^\mu dx^\nu + dz^2 + \beta(z)d\theta^2)$, where $\beta(z) = \gamma(z)/\sigma(z)$. Moreover, performing a change of the dependent variable as

$$\phi_n(z) = K(z)\Psi_n(z) , \quad K(z) = \sigma^{-1}(z)\beta^{-\frac{1}{4}}(z) , \quad (20)$$

we turn the Eq. (13) into the following Schrödinger-like equation for the $\Psi_n(z)$ function:

$$-\ddot{\Psi}_n(z) + U(z)\Psi_n(z) = m_n^2 \Psi_n(z) , \quad (21)$$

where the dots represent derivatives with respect to the z coordinate and the analogue quantum potential $U(z)$ has the form

$$U(z) = \mathcal{W}^2(z) + \dot{\mathcal{W}}(z) , \quad \mathcal{W}(z) = \left(\frac{\dot{\sigma}}{\sigma} + \frac{1}{4} \frac{\dot{\beta}}{\beta} \right) . \quad (22)$$

Note that the Eq. (21) preserves the analogue supersymmetric quantum mechanics form:

$$\left[\frac{d}{dz} + \mathcal{W}(z) \right] \left[-\frac{d}{dz} + \mathcal{W}(z) \right] \Psi(z) = \mathcal{Q}\mathcal{Q}^\dagger \Psi(z) = m^2 \Psi_m(z) . \quad (23)$$

Hence, the absence of tachyonic modes is guaranteed and the stability of the spectrum is ensured.

With the above mentioned changes of variables, the boundary conditions become

$$\frac{\dot{\Psi}(0)}{\Psi(0)} = -\frac{\dot{K}(0)}{K(0)} , \quad \frac{\dot{\Psi}(\infty)}{\Psi(\infty)} = -\frac{\dot{K}(\infty)}{K(\infty)} . \quad (24)$$

The orthogonal condition is modified to

$$\int_0^\infty \Psi_{m_i}^*(z) \Psi_{m_j}(z) dz = \delta_{ij} . \quad (25)$$

It is important to mention that the gravitational zero-mode must reproduce the four-dimensional gravity on the brane [8, 10]. Hence, the solution for $m = 0$ of the Eq. (21) is

$$\Psi_0(z) = C \sigma(z) \beta^{\frac{1}{4}}(z) , \quad (26)$$

where C is a normalization constant. For the GS model, the zero mode in the z variable is analytically evaluated as

$$\Psi_0(z) = \sqrt{\frac{24}{c^3}} \left(z + \frac{2}{c} \right)^{-2} . \quad (27)$$

It is important to mention that the gravitational zero mode is always non-singular and normalizable for all values of the resolution parameter, as showed in the Fig. 3. Moreover, the massless solution shares similar profile with the energy density due to the core peak displaced from the origin for small values of a . Such property was also observed in the string-cigar braneworld [31, 43] and in the Abelian-vortex braneworld [29]. For great values of the resolution parameter, the graviton zero mode has its peak at the origin. This is in accordance with the GS thin-string model [23], in the Bounce 6D model [46] and the Torrealba's string-vortex [30]. This aspect of resolution parameter influences the corrections to Newton's Law.




Figure 3. Gravitational massless mode in the resolved conifold braneworld. The thin string-like case is embedded.

Besides, for $m \neq 0$, the solutions of Eq. (21) in the GS scenario are given by

$$\Psi_m(z) = \sqrt{z + \frac{2}{c}} \left[C_1 J_{\frac{5}{2}}\left(mz + 2m/c\right) + C_2 Y_{\frac{5}{2}}\left(mz + 2m/c\right) \right], \quad (28)$$

where C_1 and C_2 are constants.

We will now verify if the gravitational interactions mediated by the Kaluza-Klein modes are in agreement with the four-dimensional laws of gravity. For such purpose, we will consider a minimal coupling of matter to gravity and look for the values of the coupling constants. To compute the gravitational potential between two point-like particles with masses M_1 and M_2 on the brane, we rewrite the Kaluza-Klein decomposition as

$$h_{\mu\nu}(x^\zeta, z, \theta) = \frac{1}{M_{Pl}^2} K(z) \sum_{n=0}^{\infty} \tilde{h}_{\mu\nu}^{(n)}(x^\zeta) \Psi_n(z) e^{il\theta}. \quad (29)$$

Hence, we have the Lagrangian

$$\mathcal{L}_{\mathcal{M}}(\Phi, g'_{MN}) = \mathcal{L}_{\mathcal{M}}(\Phi, g_{MN}) - \sum_{n=0}^{\infty} a_n(z) \tilde{h}_{\mu\nu}^{(n)}(x^\zeta) T^{\mu\nu}(x^\zeta), \quad (30)$$

where the expression of the gravity-matter coupling constants is given by:

$$a_n(z) = \frac{1}{2M_{Pl}^2} \frac{\Psi_n(z) e^{il\theta}}{\sigma(z) \beta^{\frac{1}{4}}(z)}. \quad (31)$$

We are now able to compute the static potential generated by the exchange of the zero-mode and massive KK states. Like in the case of a Yukawa interaction, it is given by

$$V(r) = -\frac{1}{4\pi} \sum_{n=0}^{\infty} \left| a_n(z = \bar{z}) \right|^2 \frac{e^{-m_n r}}{r}, \quad (32)$$

where $\bar{z}$ is the position of the maximum of the energy density on the z coordinate. Therefore, on the 3-brane, the gravitational potential between two point masses, M_1 and M_2 , will receive a Yukawa-like contribution from the discrete nonzero modes as:

$$V(r) \sim G_N \frac{M_1 M_2}{r} + G_N \frac{M_1 M_2}{r} \sum_{n>0}^{\infty} \frac{\Psi_n^2(\bar{z})}{\sigma^2(\bar{z}) \beta^{\frac{1}{2}}(\bar{z})} e^{-m_n r}, \quad (33)$$

Therefore, we have an expression for general thick braneworld scenario. In order to analyze the effects of the gravitational Kaluza-Klein modes in the resolved conifold scenario to the Newton's law of gravity, we need to compute the mass spectrum and the corresponding eigenfunctions using suitable numerical methods. This will be accomplished in the next section.

V. NUMERICAL ANALYSIS

In the previous sections we presented the main features of a resolved conifold braneworld model and the gravity localization in this scenario. Furthermore, a general expression for the correction in the gravitational potential due to Kaluza-Klein spectrum provided from a six-dimensional thick braneworld scenario was derived. We are now able to compute this correction due to the resolved conifold braneworld. Note that the leading quantities in Eq. (33) are the masses m_n and the wave functions $\Psi_n(z)$ which are eigensolutions of the eigenvalue problems (17) and (21). However, due to the involved form of the Eq. (17), such quantities can not be obtained analytically. Then, we have used suitable numerical method to solve the Sturm-Liouville problems.

We have used the matrix method [48] based in finite differences schemes with second order truncation error to solve the Sturm-Liouville problem (17). The matrix method is a numerical technique used to approximate the first eigenvalues and eigenfunctions of Sturm-Liouville problems. Indeed, in braneworld models, only the small masses are of physical interest, favouring us the use of the finite differences methods. This technique was used in the braneworld context to attain the massive spectrum in five [16] and six dimensions [43–45].

Here we are interested in to study the effects of the Kaluza-Klein tower in terms of the resolution parameter a , since the parameter related to the geometric flow c acts as an energy scale [43–45]. Then, we solved the Eq. (17) for different values of the resolution parameter a and holding $c = 1.0$ fixed. We discretized the domain $\rho \in [0.0, 8.0]$ with $N = 1\,600$ subdivisions, in order that the constant stepsize is obtained as $h = 0.005$.

We plot the first twenty mass eigenvalues in the Fig. 4. The spectrum is real and monotonically increasing, as expected. Note that, as the resolution parameter increases, the spectrum enhances

in magnitude and the spacing between the masses enlarges. Note further that the masses exceed the limit of $m = 1.0$ for high values of a , and belong to a transplanckian regime (remember that $m_n \ll c = 1.0$ [8, 10, 23, 24]). Thus, the resolution parameter must assume moderate values to have physically acceptable states.

With the mass eigenvalues in hands, we solve the Schrödinger-like equation (21) with the potential function (22) for the eigenvalues obtained previously. Such procedure was also followed in a sophisticated five-dimensional braneworld scenario called, hybrid brane [16]. Since the transformation of coordinates $z(\rho)$ has no analytical expression for the warp factor in the Eq. (8), a numerical approximation was necessary to construct the analogue quantum potential. After a numerical quadrature of $z(\rho)$ for the warp factor $\sigma(\rho)$, we were able to compute the functions $\sigma(z)$ and $\beta(z)$ using cubic spline interpolation. Then, a numerical approximation for $U(z)$ was possible.

We plot the analogue quantum potential $U(z)$ for the resolved conifold in the Fig. 5 for different values of a and for $c = 1.0$. The thin string-like case is embedded. In the new coordinate z , the domain is stretched to $z \in [0, 65]$. Note that the resolution parameter removes the singularity at the origin for the RC model. Such feature was also verified in the case of scalar and matter fields [32, 38, 39]. For $a = 0$, the analogue quantum potential has a singularity at the origin. Such behaviour is also found in the string-cigar model [31]. Moreover, the depth of the potential well and the height of barrier are smoothed by the resolution parameter with $a \neq 0$, which has no major changes for $a > 1.0$.

To analyze how the resolution parameter affects the gravity phenomenology of the resolved conifold braneworld, we solved the Schrödinger-like equation using the Numerov algorithm [49] with




Figure 4. Gravitational Kaluza-Klein spectrum for different values of the resolution parameter a .

the mass eigenvalues obtained previously as solution of the Sturm-Liouville problem (13). We plot in the figures 6 and 7 the numerical solutions for $a = 0$ and $a = 0.5$, respectively, for different mass eigenvalues. A noteworthy result concerns to the seventh wavefunction for the singular conifold ($a = 0$), which has a resonant profile (massive wavefunctions having large amplitude near the brane [50, 51]). The mass of this particular state is $m_7 = 0.351145$, which is in accordance with the limit that a resonant mass must be $m^2 < \max(U)$ [10, 52]. In the present case, $\sqrt{\max(U)} = 0.444009$. It is important to mention that graviton resonant states were also found in another thick string-like brane scenario with conical singularity, the string-cigar model [43]. Furthermore, the resolved case does not present resonant modes. We have solved the radial equations for several values of a .

Since the main quantities that contribute to the corrections in the gravitational potential are the exponentially suppressed Kaluza-Klein masses and the corresponding eigenfunctions – see Eq. (33) – it can be seen that the resonant state is the massive mode with largest contribution to the correction of the Newton’s law of gravitation. Interestingly, in the Ref. [53], a five dimensional asymmetric brane model exposes a resonant state performing a stronger contribution to the correction in the Newtonian potential. On the other hand, in the symmetric brane models, the first eigenstate contributes highly for such correction [10, 16].

From the aforementioned results, we are led to comprehend from the Eq. (33) that the $n = 7$ (resonant eigenstate) is the majority term of the sum. To express more clearly the behaviour of the correction in the Newtonian potential, we computed the Eq. (33) for all twenty eigensolutions $\{m_n, \Psi_n(\bar{z})\}$. It is important to mention that the position of the peak in the resonant wave




Figure 5. Analogue quantum potential for the gravitons in the Resolved Conifold braneworld scenario. The thin string-like case $U_{GS}(z)$ is embedded.

function $\bar{z}$ and the position of the maximum of the energy density in the z coordinate are very close. Thus, the graviton exchange is being computed near the point where the energy of the brane is maximum. We present this result in the Fig. 5. Note that the gravitational force between two point-like masses is slightly augmented for short distances.

VI. CONCLUSIONS AND PERSPECTIVES

In this work we have studied the gravity fluctuations in a brane placed at a transverse warped resolved conifold (RC). The RC model is a thick string-like braneworld model, whose transverse space is a 2-section of a resolved conifold. Such brane model traps the scalar [32], gauge [38] and fermionic [39] fields. In this work we show that the RC braneworld also traps the gravitational field. The RC model has a parameter a , called resolution parameter, which removes the conical singularity. This parameter has strong influences in the geometric and physical properties of the model [32, 38, 39]. We therefore, studied the effects of the resolution parameter in the phenomenological implications of the resolved conifold model via small deviations in the Newtonian potential. We derived a general expression for the correction in the gravitational potential between two point-like sources of mass due to the gravitational Kaluza-Klein (KK) states in a thick string-like braneworld. The correction has an exponentially suppressed mass term.

The massless mode, which is responsible to reproduce the four-dimensional gravity, is trapped at the brane and is non-singular and normalizable for all the values of the resolution parameter. Moreover, the massless mode is displaced from the origin sharing similar profile with the energy




Figure 6. Normalized graviton wavefunctions for the Conifold braneworld scenario ($a = 0$). The seventh eigenstate has a resonant profile.




Figure 7. Normalized graviton wavefunctions for the Resolved Conifold braneworld scenario ($a \neq 0$).

density of the brane.

Using suitable numerical methods, we attained the mass spectrum of the KK graviton. The spectrum is real and monotonically increasing, as expected. However, as the resolution parameter increases, the spectrum enhances in magnitude so that for high values of a , the masses belong to a transplanckian regime. Thus, the resolution parameter must assume moderate values to have physically acceptable states. We also computed the analogue quantum potential. The resolution parameter removes the singularity of the potential at the origin. Nonetheless, for high values of a the potential well has not significant changes.

We solved numerically the Schrödinger-like equation for the analogue quantum potential using the mass eigenvalues for several values of the resolution parameter. For $a = 0$ (singular conifold) we found a resonant mode which is the seventh eigenstate. However, for $a \neq 0$ (resolved conifold), we have not found resonant states. Since the value of the wavefunction at the core peak of the brane is the quantity which is used to compute the correction in the gravitational potential, the resonant state is the major contributor to this correction. A noteworthy result is that a highly excited state is the responsible for the correction in the Newtonian potential rather than the first eigenstate.

We present results for a fixed energy scale for the mass $m \ll c = 1.0$, such that the resolution parameter a should preferably be smaller than one to have physically acceptable states. Similar results were obtained for different values of c . Motivated by this peculiar result, we intend for future works, to study the phenomenological consequences of the resolved conifold in the context of the Configurational Entropy [54], computing bounds in the resolution parameter. Moreover, we will study the correction to Coulomb's Law [55] in thick string-like six dimensional scenarios.

VII. ACKNOWLEDGMENTS

The authors thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPQ), and Fundação Cearense de apoio ao Desenvolvimento Científico e Tecnológico (FUNCAP) for financial support.

-
- [1] P. Horava and E. Witten, Nucl. Phys. B **460**, 506 (1996).
 - [2] N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **429**, 263 (1998);
I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, Phys. Lett. B **436**, 257 (1998).

- [3] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999).
- [4] N. Arkani-Hamed, S. Dimopoulos and J. March-Russel, Phys. Rev. D **63**, 064020 (2001).
- [5] N. Arkani-Hamed, S. Dimopoulos, G. Dvali and N. Kaloper, J. High Energy Phys. 012 (2000) 010.
- [6] J. W. Chen, M. A. Luty and E. Pontón, J. High Energy Phys. 09 (2000) 012.
- [7] J. Khoury, B. A. Ovrut, P. J. Steinhardt and N. Turok, Phys. Rev. D **64**, 123522 (2001);
S. Räsänen, Nucl. Phys. B **626**, 183 (2002).
- [8] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999).
- [9] W. D. Goldberger and M. B. Wise, Phys. Rev. Lett. **83**, 4922 (1999).
- [10] C. Csáki, J. Erlich, T. J. Hollowood and Y. Shirman, Nucl. Phys. B **581**, 309 (2000).
- [11] M. Gremm, Phys. Lett. B **478**, 434 (2000).
- [12] D. Bazeia and A. R. Gomes, J. High Energy Phys. 05 (2004) 012.
- [13] V. Dzhunushaliev, V. Folomeev and M. Minamitsuji, Rep. Prog. Phys. **73**, 066901 (2010).
- [14] G. Germán, A. Herrera-Aguilar, A. M. Kuerten, D. Malagon-Morejon and R. da Rocha, J. Cosmol. Astropart. Phys. 01 (2016) 047.
- [15] W. T. Cruz, R. V. Maluf, L. J. S. Sousa and C. A. S. Almeida, Annals Phys. **364**, 25 (2016);
W. T. Cruz, R. V. Maluf, D. M. Dantas and C. A. S. Almeida, arXiv:1512.07890 [hep-th].
- [16] D. F. S. Veras, W. T. Cruz, R. V. Maluf and C. A. S. Almeida, Phys. Lett. B **754**, 201 (2016).
- [17] W. D. Goldberger and M. B. Wise, Phys. Rev. D **60**, 107505 (1999).
- [18] S. J. Huber and Q. Shafi, Phys. Lett. B **498**, 256 (2001).
- [19] A. Kehagias and K. Tamvakis, Phys. Lett. B **504**, 38 (2001).
- [20] A. G. Cohen and D. B. Kaplan, Phys. Lett. B **470**, 52 (1999).
- [21] R. Gregory, Phys. Rev. Lett. **84**, 2564 (2000).
- [22] I. Olasagasti and A. Vilenkin, Phys. Rev. D **62**, 044014 (2000).
- [23] T. Gherghetta and M. E. Shaposhnikov, Phys. Rev. Lett. **85**, 240 (2000).
- [24] I. Oda, Phys. Lett. B **496**, 113 (2000).
- [25] Y. -X. Liu, L. Zhao and Y. -S. Duan, J. High Energy Phys. 04 (2007) 097.
- [26] V. Santos and C. A. S. Almeida, Phys. Lett. B **718**, 1114 (2013).
- [27] T. G. Rizzo, J. High Energy Phys. **11** (2010) 156.
- [28] P. Tinyakov and K. Zuleta, Phys. Rev. D **64**, 025022 (2001).
- [29] M. Giovannini, H. Meyer and M. E. Shaposhnikov, Nucl. Phys. B **619**, 615 (2001).
- [30] R. S. Torrealba, Gen. Rel. Grav. **42**, 1831 (2010).
- [31] J. E. G. Silva, V. Santos and C. A. S. Almeida, Class. Quant. Grav. **30**, 025005 (2013).
- [32] J. E. G. Silva and C. A. S. Almeida, Phys. Rev. D **84**, 085027 (2011).
- [33] J. C. B. Araújo, J. E. G. Silva, D. F. S. Veras and C. A. S. Almeida, Eur. Phys. J. C **75**, 127 (2015).
- [34] J. Garriga and M. Porrati, J. High Energy Phys. 08 (2004) 028.
- [35] M. Gogberashvili, P. Midodashvili and D. Singleton, J. High Energy Phys. 08 (2007) 033.
- [36] S. Aguilar and D. Singleton, Phys. Rev. D **73**, 085007 (2006).

- [37] J. M. Frère, M. Libanov, S. Mollet and S. Troitsky, J. High Energy Phys. 08 (2013) 078.
- [38] F. W. V. Costa, J. E. G. Silva and C. A. S. Almeida, Phys. Rev. D **87**, 125010 (2013).
- [39] D. M. Dantas, J. E. G. Silva and C. A. S. Almeida, Phys. Lett. B **725**, 425 (2013).
- [40] I. Navarro and J. Santiago, J. High Energy Phys. 02 (2005) 007.
- [41] P. Candelas and X. C de la Ossa, Nucl. Phys. B **342**, 246 (1990).
- [42] L. A. P. Zayas and A. A. Tseytlin, J. High Energy Phys. 11 (2000) 028.
- [43] D. F. S. Veras, J. E. G. Silva, W. T. Cruz and C. A. S. Almeida, Phys. Rev. D **91**, 065031 (2015).
- [44] F. W. V. Costa, J. E. G. Silva, D. F. S. Veras and C. A. S. Almeida, Phys. Lett. B **747**, 517 (2015).
- [45] D. M. Dantas, D. F. S. Veras, J. E. G. Silva and C. A. S. Almeida, Phys. Rev. D **92**, 104007 (2015).
- [46] L. J. S. Sousa, C. A. S. Silva, D. M. Dantas and C. A. S. Almeida, Phys. Lett. B **731**, 64 (2014).
- [47] D. M. Dantas, R. da Rocha and C. A. S. Almeida, arXiv:1512.07888 [hep-th].
- [48] P. Amodio and G. Settanni, J. Numer. Anal. Indust. Appl. Math. **6**, 1 (2011).
- [49] B. V. Numerov, Mon. Not. R. Astron. Soc. **84**, 592 (1924);
B. V. Numerov, Astron. Nachr. **230**, 359 (1927).
- [50] C. A. S. Almeida, M. M. Ferreira, Jr., A. R. Gomes and R. Casana, Phys. Rev. D **79**, 125022 (2009).
- [51] Y. Liu, J. Yang, Z. Zhao, C. Fu, and Y. Duan, Phys. Rev. D **80**, 065019 (2009).
- [52] C. Csáki, J. Erlich, and T. J. Hollowood, Phys. Rev. Lett. **84**, 5932 (2000).
- [53] G. Gabadadze, L. Grisa and Y. Shang, J. High Energy Phys. 08 (2016) 033;
A. Araujo, R. Guerrero and R. O. Rodriguez, Phys. Rev. D **83**, 124049 (2011).
- [54] R. A. C. Correa, D. M. Dantas, C. A. S. Almeida and R. da Rocha, Phys. Lett. B **755**, 358 (2016);
R. A. C. Correa, D. M. Dantas, C. A. S. Almeida and P. H. R. S. Moraes, arXiv:1607.01710 [hep-th].
- [55] H. Guo, A. Herrera-Aguilar, Y. X. Liu, D. Malagon-Morejon and R. R. Mora-Luna, Phys. Rev. D **87**, 095011 (2013);
R. Cartas-Fuentevilla, A. Escalante, G. Germn, A. Herrera-Aguilar and R. R. Mora-Luna, J. Cosmol. Astropart. Phys. 05 (2016) 026.